



## Phylogenetic assay and pollen structure of few species of the genus *amaranthus* L.

Pinkie Cherian\*, D. Sheela and Durga K.V.

Department of Botany, St.Teresa's College, Ernakulam, Kerala-682011, India  
pinkie.cherian@yahoo.co.in

Available online at: [www.isca.in](http://www.isca.in), [www.isca.me](http://www.isca.me)

Received 22<sup>nd</sup> March 2017, revised 23<sup>rd</sup> May 2017, accepted 8<sup>th</sup> June 2017

### Abstract

In *Amaranthaceae*, morphology based identification methods are usually time consuming and may sometimes lead to misidentification and always may not provide good resolution at the species levels. The phenotypic variability of the taxa may lead to misidentifications and creation of new false identity. DNA sequencing has been used to explain evolutionary relationships for more than 20 years in molecular systematics. The aims of DNA barcoding include identification of known specimens/species and new discovery of unknown plant species for enhancing taxonomy for the good of the science and betterment of society. The study basically emphasised on palynological studies and molecular profiling of *Amaranthus* species using universal markers *rbcL* and *matK*. The pollen morphology of the species of *Amaranthus* shows significant differences in polar length and equatorial diameter whereas the aperture was pantoporate with evenly distributed microspines. The phylogenetic assay showed bootstrap value of 96 and 98 for *matK* while 79 and 98 for *rbcL* dendrograms. All barcodes yield quality sequences.

**Keywords:** DNA Barcoding, *rbcL*, *matK*, Palynology.

### Introduction

The flowering plant family *Amaranthaceae* which corresponds to the classical family *Amaranthaceae* Juss. 69 genera and 772 species as well as the twice larger related family *Chenopodiaceae* have been subject to repeated taxonomical revisions from the time they were first described (1789 and 1799) respectively to the present<sup>1-3</sup>. Recently it has been proposed to combine them into one large family *Amaranthaceae* as a result of molecular analysis<sup>4</sup>. The morphology of pollen grains, is an important source of information for plant systematics. In certain cases, palynological data have been crucial for taxonomic conclusions<sup>5</sup>. But for closely related species of the genus *Amaranthus* L., such data can lead to confusions regarding the classification though pollen grains morphological studies shows very slight variation. Traditionally, most plant identifications are based on morphological characters, but such identification is not always reliable and efficient<sup>6</sup>.

DNA barcoding basically relies on short and standardized gene regions for the identification of plant species. The agricultural and horticultural applications of barcoding such as for marketplace regulation and copyright protection remain poorly explored. This study examines the use of effectiveness/quality of the standard plant barcode markers (*matK* and *rbcL*) for the identification of plant species that are medicinal and are of interest for pharmaceutical industry both in private and public nurseries and to authenticate the medicinal plants. A wide variety and different range of molecular techniques including

the random amplified polymorphic DNA (RAPD), amplified fragment length polymorphism in plant (AFLP), restriction fragment length polymorphism of the selected taxa (RFLP), microsatellite and single nucleotide polymorphism (SNP) have been proposed to identify plant species/specimen and cultivars<sup>7-11</sup>. DNA barcoding method has emerged as a relatively new/novel and perhaps more universal tools with which to analyze diversity of both plants and animals and to fix specimens to their respective species even in the absence/unavailability of key morphological diagnostic features<sup>12,13</sup>. Although there are still some reserves against the performance of DNA barcoding as compared, for example, to morphology, an early study, through a thorough comparison of DNA barcoding and morphology-based species identification recorded a number of limitations to the morphology particularly when it comes to cryptic species<sup>2</sup>. As a taxonomic tool this technique has widely acceptable and also has been successfully used in large scale biodiversity projects where regional flora and fauna are documented including regulated and threatened taxa<sup>14-17</sup>. Although a number of plant loci including, *trnH-psbA*, *rpoc1*, *rpoB*, *trnL*, *rbcL* and *matK* were initially proposed as potential plant barcodes based on assessments of similarity with genebank and recoverability, sequence derived quality and levels of plant species discrimination, the Consortium for the Barcode of Life recommended the 2-loci combination of *rbcL* + *matK* as the most standard plant barcode for analysis<sup>18-23</sup>.

The DNA barcode data generated in the present study will serve in the future in commercial agricultural and medicinal plant

industries for the purpose of control of counterfeited product, and could also serve in ecological studies of local flora as demonstrated elsewhere<sup>24-26</sup>. In addition, the evolutionary processes such as hybridization especially interspecific hybridisation and different polyploidy are common in plants, so such species boundaries are difficult to define<sup>27,28</sup>. Thus, screening for single or multiple regions using appropriate primers are necessary for DNA barcoding studies in the nuclear and plastid genomes in plants that has been an important point of research. Since *matK* and *rbcL* sequences from Amaranthaceae were the most abundant in GeneBank, they were chosen for the study. Similarly authentication is a critical aspect of research in selection of plants for study, so an attempt was done to authenticate the *Amaranthus* L. using the palynological and DNA barcodes data in order to make a unique identity among the plants.

### Materials and methods

The fresh leaves of *Amaranthus spinosus* L., *Amaranthus caudatus* L., *Amaranthus tricolor* L., *Amaranthus dubius* Mart., and *Amaranthus viridis* L. were used for isolating genomic DNA. Pollen grains were collected from the anther of 15 flowers of each species and fixed in glacial acetic acid for acetolysis as per the technique<sup>29</sup>. The sculpturing pattern, values of P (pollen axis ratio) and E (equatorial diameter) and aperture number were viewed and data were measured using the Scanning Electron Microscope (SEM) JSM-6390 LA coated with gold, examined and photographed using JSM-6390 LA. All the experiments were evaluated statistically with SPSS version 20.0, the results were represented in mean±SEM (standard error of mean). One way analysis of variance (ANOVA) followed by DMRT to find out any significant difference in pollen characters among five *Amaranthus* sps. resulting from analytical experiments carried out. P value less than 0.05 were adopted as statistically different.

The DNA was isolated by using branded Sigma kit GenElute Plant Genomic DNA Mini-preparation Kit. GeneAmp PCR System 9700, Applied Biosystems, a PCR thermal cycler for PCR amplification, using the primers of *rbcL* and *matK*. The primer details were given in table 1 and PCR amplification data for conditions provided are given in Table-2.

**Table-1:** The universal primers *rbcL* and *matK* and their sequences.

Target gene	Primer	Direction	Sequence strand (5' → 3')
<i>matK</i>	390 f	forward	CGATCTATTCATTCAA TATTTTC
	1326r	reverse	TCTAGCACACGAAAGT CGAAGT
<i>rbcL</i>	rbcLa_f	Forward	ATGTCACCACAAACA GAGACTAAAGC
	rbcL724_rev	Reverse	GTAATAATCAAGTCCAC CRCG

**Table-2:** PCR amplification profile.

<i>matK</i>		<i>rbcL</i>	
98 °C	-30 sec	98 °C	-30 sec
98 °C	- 5 sec	98 °C	- 5 sec
50 °C	-10 sec	60 °C	-10 sec
72 °C	-15 sec	72 °C	-15 sec
72 °C	- 60 sec	72 °C	- 60 sec
4 °C	- ∞	4 °C	-∞


Sequencing reaction was done in a PCR thermal cycler named GeneAmp PCR System 9700, Applied Biosystems) using the BigDye Terminator v3.1 Cycle sequencing Kit of Applied Biosystems, USA model. The sequence quality was checked Applied Biosystems Sequence Scanner Software v1. Using Geneious Pro v5.6 Sequence alignment and required editing of the obtained sequences were carried out<sup>30</sup>. The DNA sequences of *Amaranthus* spp. under study were subjected to BLAST analysis for better identification at the species level. Sequences obtained were aligned and compared using Multiple Sequence Alignment software program of BioEdit Sequence Alignment Editor, CLUSTAL W Multiple Alignment<sup>31,32</sup>. DNA barcodes namely *rbcL* and *matK* for constructing neighbouring tree model using MEGA 7.0, and a tree was constructed using a combination of *rbcL* and *matK*<sup>33</sup>.

### Results and discussion

The SEM photomicrographs of radial, longitudinal axis, pore aperture, aperture number and distance are shown in Figure-1. The pore structure is similar to the Type II of that is the pores possess microspines, granulate surface and are evenly spread, with numerous pores<sup>34</sup>. The present palynological result support existing data which have indicated that *Amaranthus* is pantoporate in terms of pore number. Most of the species examined in Amaranthaceae have pantoporate with *Amaranthus* type of pollen<sup>35</sup>. The aperture of the pollen examined showed great variation in *Amaranthus tricolor* L., where their size and number is reduced as reported that there is a *Amaranthus* type of porate aperture in the family Amaranthaceae<sup>36</sup>. The pollen grains of *Amaranthus* species are spheroidal to oblate spheroidal, pantoporate and pantotreme, with this a key is prepared in order to make a comparison and identification among the species.

Shape- Spheroidal  
 Diameter 22.12-22.21µm  
 Aperture distance 3.23 -3.66 µm ..... *Amaranthus viridis* L.  
 Shape- Oblate-spheroidal  
 Diameter 21.80-21.84µm  
 Aperture distance 1.00 -1.22 µm ..... *Amaranthus spinosus* L.  
 Diameter 18.00-19.86 µm  
 Aperture distance 2.05-2.57 µm ..... *Amaranthus tricolor* L.

Shape- Prolate-spheroidal  
 Diameter 20.32-21.34µm  
 Aperture distance 3.02-3.14 µm .....*Amarantus dubius* Mart.  
 Diameter 20.46 -20.83 µm  
 Aperture distance 2.12-2.21 µm ..... *Amarantus caudatus* L.



**Figure-1:** SEM photograph of *Amaranthus* L. pollen.

This study demonstrated differences in pollen characteristics among the *Amaranthus* species. However, pollen morphology/palynological data will have limited use in species identification because of similarities across the species. To

increase the usefulness of pollen morphology in species identification, additional analysis of naturally occurring population as well as sampling from different geographical regions would be needed to account for the study<sup>37</sup>. The statistical analysis of pollen characters using SPSS version 20.0 by Duncan's Multiple Range test ( $\alpha/p = 0.05$ ) showed that there is a significant difference between the species of *Amaranthus* L. as because the  $p < 0.05$ . Thus it infers that the palynological characters are significant among five selected species. Similar attempt was made to pollen morphological differences in *Amaranthus* L. species and the hybrids formed by interspecific hybridisation where the mean value with same letter are not significant as reported<sup>37</sup>.

From the DNA barcoding studies using the universal primers *rbcl* and *matK*, the study revealed the correct differentiation among *Amaranthus* species. It has been proved experimentally that DNA markers can act as a powerful/authentic tool for identification of cultivars and species for phylogenetic evaluation<sup>38,39</sup>.

The accession numbers of the DNA sequences submitted and size of the sequences are given in the Table-5. The tree constructed by neighbour joining tree feature using MEGA 7 is given in the Figure-2.

**Table-4:** Pollen characters among 5 *Amaranthus* species

Species	<i>Amarantus viridis</i> L.	<i>Amarantus spinosus</i> L.	<i>Amarantus dubius</i> Mart.	<i>Amarantus caudatus</i> L.	<i>Amarantus tricolor</i> L.
Pore distance	1.254 <sup>d</sup> ±0.006	1.19 <sup>d</sup> ±0.022	1.172 <sup>s</sup> ±0.19	1.226 <sup>d</sup> ±0.01	1.146 <sup>c</sup> ±0.02
Aperture distance	3.448 <sup>c</sup> ±0.081	2.924 <sup>c</sup> ±0.146	2.672 <sup>d</sup> ±0.07	2.73 <sup>c</sup> ±0.06	2.534 <sup>d</sup> ±0.07
Diameter(l)	20.872 <sup>b</sup> ±0.85	20.634 <sup>b</sup> ±0.766	19.91 <sup>c</sup> ±0.48	20.006 <sup>b</sup> ±0.49	19.652 <sup>c</sup> ±0.45
Diameter(e)	21.086 <sup>b</sup> ±0.321	21.376 <sup>b</sup> ±0.257	21.308 <sup>b</sup> ±0.32	21.386 <sup>b</sup> ±0.36	21.364 <sup>b</sup> ±0.37
Aperture number	24.2 <sup>a</sup> ±0.74	25.8 <sup>a</sup> ±0.374	28 <sup>a</sup> ±0.44	30 <sup>a</sup> ±0.89	16.8 <sup>a</sup> ±0.58

Each value that is expressed in as mean±Std. Error done in triplicates. Data analysed by SPSS version 20.0 by Duncan's Multiple Range test ( $\alpha/p=0.05$ ). Mean values followed by the different Superscript in the columns are significantly different among the pollen characters.


**Table-5:** Accession number of sequence submitted in the GenBank.

Plant samples	Place of Collection	rbcl Accession No.	matK Accession No.
<i>A. viridis</i>	Edakochi	KJ773261	KJ772535
<i>A. spinosus</i>	Thevara	EF590496	EF590394
<i>A. dubius</i>	Thrissur	KX090210	KX090202
<i>A. caudatus</i>	Boat jetty	KX090209	KC747133
<i>A. tricolor</i>	Kodugallur	JF940812	JF953165

**Table-6:** Sequence data from GenBank for 5 medicinal plants.

Species	Gene	%of match with Gen Bank
<i>A. viridis</i>	<i>matK</i> <i>rbcL</i>	99% with 4 species of <i>Amaranthus</i> 99-97% to other <i>Amaranthus</i> spp.
<i>A. spinosus</i>	<i>matK</i> <i>rbcL</i>	99% match with 3 species 100% similar
<i>A. dubius</i>	<i>matK</i> <i>rbcL</i>	100-99% similar 100-99% to multiple <i>Amaranthus</i> spp.
<i>A. caudatus</i>	<i>matK</i> <i>rbcL</i>	98% to 4 <i>Amaranthus</i> spp. 100-99% to multiple <i>Amaranthus</i> spp.
<i>A. tricolor</i>	<i>matK</i> <i>rbcL</i>	100% to 3 <i>Amaranthus</i> spp. 100% with multiple <i>Amaranthus</i> spp.

% match shows how closely the barcode sequences matched with the other accessions in GenBank for *Amaranthus* L.



**Figure-2:** Phylogenetic tree constructed using two universal barcodes.

The currently available DNA sequences in the GenBank demonstrate the different challenge of discrimination power among the species. Table-5 lists out the barcoding sequences of five pharmaceutically important medicinal plants and indicates how well each sequence matches or identifies the selected plant. The results can vary by species and by gene region or target site. For example, using the *rbcL* shows good percentage of similarity with some *Amaranthus* species than *matK*. Both *matK* and *rbcL* nucleotide sequences had been previously successful for determination of phylogenetic relationships among taxa of other angiosperm plants<sup>40-42</sup>.

Using the Neighbor-Joining method the evolutionary history was inferred<sup>43</sup>. As with the case of tree constructed using *matK*, the sum of branch length = 0.01871936 is shown in the optimal tree. While tree constructed using *rbcL* primer showed branch length of 0.00720922. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (with

100 replicates) are shown next to the selected branches<sup>44</sup>. The tree is drawn to scale, with branch lengths in the same units as those of the evolutionary distances that are used to infer the phylogenetic tree constructed. The analysis basically involved 5 nucleotide sequences and all positions containing gaps and missing data were avoided. Evolutionary analyses were conducted in MEGA7<sup>33</sup>.

The robustness of the trees and branch support was estimated by bootstrap analysis. In the present study the phylogenetic assay showed bootstrap value of 96 and 98 for *matK* while 79 and 98 for *rbcL* dendrograms. The boot strap value was found to be more for *matK* than *rbcL* primer. The *matK* sequence provide good resolution within many angiosperm orders from earlier reports. Combination or combined analysis of *matK* and other rapidly evolving DNA regions with available multi-gene data sets will have the strong potential to enhance the resolution and internal support in deep level angiosperm phylogenetics and

provide additional insights into angiosperm evolution<sup>42</sup>. Similarly from the dendrogram of *rbcL* and *matK*, *A. spinosus* and *A. dubius*, *A. tricolor* and *A. viridis* showed same clade and *A. caudatus* showed distinct clade as shown in Figure-2. Thus use of these molecular markers in addition to the classical methods (palynology/morphology) provides more positive identification technique to locate new varieties and both can be used to authenticate the plants among the species level in more scientific way for providing better outcome.

## Conclusion

The palynological data as well as molecular data using *rbcL* and *matK* sequences shows that certain taxa *Amaranthus* L. show similarly and dissimilarly though they are morphologically similar. Constructing phylogeny tree using bioinformatics tool serve as a useful source of new information to both complement and evaluate morphological methods and as an aid to identifying those traditional morphological features that are taxonomically significant. Also from the findings the medicinal plants of pharmacologically important can be uniquely identified using DNA barcoding and provide quality control and standardization of the plant material supplied to the pharmaceutical industry for preparation of new drug formulation. Such identification is useful for avoiding the entry of different adulterants and substitutes for the preparation of Ayurvedic medicinal formulations.

## Acknowledgement

All the authors show a deep felt of gratitude to RGCB, Thiruvananthapuram for barcoding services and also MANF (2014-2015) for providing financial support for undergoing the work.

## References

1. Townsend C.C. (1993). *Amaranthaceae*. The families and genera of vascular plants. II Flowering plants. Dicotyledones. Springer Verlag, Berlin, 3(2), 70-91.
2. de Jussieu A.L. (1789). *Genera Plantarum. secundum ordines naturales disposita juxta methodum in Horto Regio Parisiensi exaratum*. Paris: Apud Viduam Herissant et Theophilum Barrois, 498.
3. Ventenat E.P. (1799). *Tableau Vegetal selon la Methode de Jussieu*. 2, 253.
4. Angiosperm Phylogeny Group (2003). An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants. *Bot. J. Linnean Soc.*, 141(4), 399-436.
5. Townsend C.C. (1974). Notes on *Amaranthaceae*. *Kew Bulletin*, 29(3), 461-475.
6. Ali M.A., Gyulai G., Hidvégi N., Kerti B., Al Hemaïd F.M.A., Pandey A.K. and Lee J. (2014). The changing epitome of species identification—DNA barcoding. *Saudi J. Biolog. Sci.*, 21(3), 204-231.
7. Keil M. and Griffin A.R. (1994). Use of random amplified polymorphic DNA (RAPD) markers in the discrimination and verification of genotypes in *Eucalyptus*. *Theor. Appl. Genet.*, 89(4), 442-450.
8. McKinnon G.E., Vaillancourt R.E., Steane D.A. and Potts B.M. (2008). An AFLP marker approach to lower-level systematics in *Eucalyptus* (Myrtaceae). *Am J Bot.*, 95(3), 368-380.
9. Besnard G., Khadari B., Villemur P. and Bervillé A. (2000). Cytoplasmic male sterility in the olive (*Olea europaea* L.). *Theor. Appl. Genet.*, 100(7), 1018-1024.
10. Ochieng J.W., Steane D.A., Ladiges P.Y., Baverstock P.R., Henry R.J. and Shepherd M. (2007). Microsatellites retain phylogenetic signals across genera in *Eucalyptus* (Myrtaceae). *Genet Mol Biol.*, 30(4), 1125-1134.
11. Ganai M.W., Polley A., Graner E.M., Plieske J., Wieseke R., Luerssen H. and Durstewitz G. (2012). Large SNP arrays for genotyping in crop plants. *Bioscience*, 37(5), 821-828.
12. Hebert P.D.N., Cywinska A., Ball S.L. and de Waard J.R. (2003). Biological identifications through DNA barcodes. *Proc. R. Soc. Lond. [Biol.]*, 270, 313-321.
13. CBOL (2009). Plant Working Group. A DNA barcode for land plants. *Proc. Natl. Acad. Sci.*, 106(31), 12794-12797.
14. Casiraghi M., Labra M., Ferri E., Galimberti A. and De Mattia F. (2010). DNA barcoding: a six-question tour to improve users' awareness about the method. *Brief. Bioinformatics*, 11, 440-453.
15. Yessoufou K., Davies J.T., Maurin O., Kuzmina M., Schaefer H., Van der Bank M. and Savolainen V. (2013). Large herbivores favour species diversity but have mixed impacts on phylogenetic community structure in an African savanna ecosystem. *Ecology*, 101(3), 614-625.
16. Maurin O., Davies T.J., Burrows J.E., Daru B.H., Yessoufou K., Muasya M.A., Van der Bank M. and Bond W. (2014). Savanna fire and the origins of "underground forests" of Africa. *New Phytologist*, 204, 201-214.
17. Janzen D.H., Hallwachs W., Blandin P., Burns J.M., Cadiou J.M., Chacon I. and Dapkey T. (2009). Integration of DNA barcoding into an ongoing inventory of complex tropical biodiversity. *Mol. Ecol. Resour.*, 9, 1-26.
18. Kress W.J., Wurdack K.J., Zimmer E.A., Weigt L.A. and Janzen D.H. (2005). Use of DNA barcodes to identify flowering plants. *Proc. Natl. Acad. Sci.*, 102(23), 8369-8374.
19. Chase M.W., Cowan R.S., Hollingsworth P.M., van den Berg C., Madriñán S. and Petersen G. (2007). A proposal

- for a standardized protocol to barcode all land plants. *Taxon.*, 56(2), 295-299.
20. Taberlet P., Coissac E., Pompanon F., Gielly L., Miquel C. and Valentini A. (2007). Power and limitations of the chloroplast *trnL*(UAA) intron for plant DNA barcoding. *Nucleic Acids Res.*, 35(3).
  21. Kress W.J. and Erickson D.L. (2007). A two-locus global DNA barcode for land plants: the coding *rbcL* gene complements the non-coding *trnH-psbA* spacer region. *PLoS ONE.*, 2(6), e508.
  22. China Plant BOL Group (2011). Comparative analysis of a large dataset indicates that ITS should be incorporated into the core barcode for seed plants. *Proc. Natl. Acad. Sci.*, 108(49), 19641-19646.
  23. CBOL Plant Working Group(2009). A DNA barcode for land plants. *Proc. Natl. Acad. Sci.*, 106(31), 12794-12797.
  24. Hoveka L.N., van der Bank M., Boatwright J.S., Bezeng B.S. and Yessoufou K. (2016). The noncoding *trnH-psbA* spacer, as an effective DNA barcode for aquatic freshwater plants, reveals prohibited invasive species in aquarium trade in South Africa. *S. Afr. J. Bot.*, 102, 208-216.
  25. Ashfaq M., Asif M., Anjum Z.I. and Zafar Y. (2013). Evaluating the capacity of plant DNA barcodes to discriminate species of cotton (Gossypium: Malvaceae). *Mol Ecol Resour.*, 13(4), 573-582.
  26. Levin R.A., Wagner W.L., Hoch P.C., Nepokroeff M., Pires J.C., Zimmer E.A. and Sytsma K.J. (2003). Family-level relationships of Onagraceae based on chloroplast *rbcL* and *ndhF* data. *Am. J. Bot.*, 90, 107-115.
  27. Rieseberg L.H., Troy E.W. and Eric J.B. (2006). The nature of plant species. *Nature*, 440, 524-527.
  28. Fazekas A.J., Kesanakurti P.R., Burgess K.S., Percy D.M., Graham S.W. and Barrett S.C.H. (2009). Are plant species inherently harder to discriminate than animal species using DNA barcoding markers?. *Molecular Ecology Resources*, 9,130-139.
  29. Erdtman G. (1960). The acetolysis method. *Svensk bot. tidskr.*, 54, 561.
  30. Drummond A.J., Ashton B., Buxton S., Cheung M., Cooper A., Duran C., Field M., Heled J., Kearse M., Markowitz S., Moir R., Stones-Havas S., Sturrock S., Thierer T. and Wilson A. (2011). Geneious Pro v5. 5.6. *View Article PubMed/ NCBI Google Scholar.*
  31. Hall T.A. (1999). BioEdit: a user- friendly biological sequence alignment editor and analysis program for windows 95/98/NT. *Nucl. Acids. Symp. Ser.*, 41, 95-98.
  32. Thompson J.D., Desmond G.H. and Toby J.G. (1994). CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Research.*, 22, 4673-4680.
  33. Kumar S., Stecher G. and Tamura K. (2016). MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution*, 33(7), 1870-1874.
  34. Borsch T. (1998). Pollen types in the Amarantaceae. Morphology and evolutionary significance. *Grana*, 37(3), 129-142.
  35. Erdtman G. (1952). Pollen Morphology and Taxonomy. Angiosperms. Almqvist and Wiksell, Stockholm.
  36. Roland F. (1971). The detailed structure and ultra-structure of an acalymmate tetrad. *Grana.*, 11, 41-44.
  37. Franssen Aaron S., Skinner Daniel Z., Al-Khatib Kassim and Horak Michael J. (2001). Pollen morphological differences in *Amaranthus* species and interspecific hybrids. *Weed Science*, 49(6), 732-737.
  38. Prevost A. and Wilkinson M.J. (1999). A new system of comparing PCR primer applied to ISSR fingerprinting of potato cultivars. *Theor. Appl. Genet.*, 98, 107-112.
  39. Wang G., Mahalingam R. and Knap H.T. (1998). C-A and G-A anchored simple sequence repeats (ASSRS) generated polymorphism in Soyabean, *Glycine max*(L.) Merr. *Theor. Appl. Genet.*, 96(8), 1086-1096.
  40. Muller K. and Borsch T. (2005). Phylogenetics of Amaranthaceae based on *matk/trnk* sequence data: evidence from parsimony, likelihood and bayesian analyses. *Annals of the Missouri Botanical Garden.*, 92(1), 66-102.
  41. Stoeckle M. (2003). Taxonomy, DNA, and the Bar Code of Life. *BioScience.*, 53(9), 796-797.
  42. Soltis D.E., Savolainen V. and Chatrou L. (2003). Angiosperm phylogeny based on *mat K* sequence information. *Am. J. Bot.*, 90(12), 1758-1776.
  43. Saitou N. and Nei M. (1987). The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution.*, 4(4), 406-425.
  44. Felsenstein J. (1985). Confidence limits on phylogenies: An approach using the bootstrap. *Evolution.*, 39, 783-791.