



Evaluation of the cytotoxicity potential of ethanolic extract of *Chrysophyllum cainito* using the brine shrimp lethality bioassay

Leonel P. Lumogdang^{1*}, Lynde Q. Bullong¹, Olga M. Nuñez² and Mylene M. Uy²

¹Department of Marine Biology, Southern Philippines of Agri-Business and Marine and Aquatic School of Technology (SPAMAST), Poblacion Malita, Davao Occidental Philippines

¹College of Arts and Sciences, Xavier University-Ateneo De Cagayan, Cagayan de-Oro Philippines

²Department of Biological Sciences, Mindanao State University-Iligan Institute of Technology (MSU-IIT), Tibanga, Iligan City Philippines

²Department of Chemistry, Mindanao State University-Iligan Institute of Technology (MSU-IIT), Tibanga, Iligan City Philippines
leonel.lumogdang@spamast.edu.ph

Available online at: www.isca.in, www.isca.me

Received 22nd July 2020, revised 13th December 2020, accepted 1st January 2021

Abstract

The Philippines has rich floral biodiversity accompanied with an abundant source of medicinal plants easily accessible in the locality. In terms of ethno-medical properties, *Chrysophyllum cainito* has been used to treat various diseases. In this study, *C. cainito* leaves were collected and evaluated for cytotoxicity using the Brine Shrimp Lethality Bioassay. The *C. cainito* leaves were extracted with water, 50:50 ethanol-water, and absolute ethanol to produce the decoction, hydro-ethanolic, and ethanolic extracts, respectively. Four concentrations (10, 100, 500, 1000 µg/ml) of the extracts were prepared and tested. The mortality rates of the brine shrimp were observed after 6 and 24 hours. The results showed that all the prepared extracts exhibited active biological activities with the ethanolic and hydro-ethanolic extracts exhibiting greater activities compared to the decoction. The ethanolic and hydro-ethanolic extracts showed toxicity effects after 24-h exposures with LC₅₀ values of 25.85 µg/ml and 84.14 µg/ml, respectively. The results indicate that the use of absolute ethanol and 50:50 ethanol-water may have successfully extracted the bioactive compounds in the *C. cainito* that have acted on the brine shrimp. The presence of active components in the extracts indicated the potential of *C. cainito* as an alternative medicine and hence requires further tests to qualitatively identify the bioactive compounds.

Keywords: Alternative medicine, bioactive, ethno-medical, extraction, mortality rate.

Introduction

The emergence of new diseases calls for new antidote that is effective, safe, and easily accessible for immediate treatments. Most populations around the world are highly dependent on traditional medicine such as the usage of herbal plants for primary health care needs¹. In fact, in developing countries such as the Philippines particularly people, who live in rural areas, resort to traditional healer when they get sick². Previous works reported the presence of chemical compounds and biological activities of plants which have yielded novel compounds with therapeutic agents which further support the viability of plants as an alternative medicine³.

One of the locally known plants which has medical importance but poorly explored scientifically is the *Chrysophyllum cainito* Linnaeus commonly known as star apple or "cainito" depending on its location⁴. *C. cainito* Linn is a tree mostly found in the tropics including Southeast Asia⁵. This tropical tree is affiliated to the family sapotaceae and is homegrown to the places of West Indies and Greater Antilles^{6,7}. *C. cainito* tree has a wide spreading crown, reaching to a height of 15 meters. Branches are usually few and slender, the young tips are conspicuously copper-colored and hid with appraised hairs.

The leaves of *C. cainito* can be characterized as leathery ovate or oblong with an estimated 7.5 to 13 centimeters long appeared with a pointed tip, edgeless or rounded at the base and has covered underneath with golden-brown, silky and soft hairs⁸.

C. cainito has several health benefits and has been chemically tested and proven⁴. Various studies were conducted through extractions of its fruit, pulp, seeds, and leaves. In 2002, polyphenolic antioxidants were isolated in the *C. cainito* fruit⁶. Fruit extracts were tested not only for their antioxidant activity but also for their antidiabetic effect⁹ and gastroprotective activity¹⁰. Moreover, a study tested the pulp and seed extracts for antimicrobial activity⁷. The tests have proven that *C. cainito* pulp and seed extracts also have great potential as antimicrobial agent for treating enteric bacterial infections and other selected pathogens. In addition, various studies also reported on the extraction of the leaves which lead to successful evaluations of the antidiabetic activity¹¹, anti-inflammatory, anti-hypersensitivity effects¹², and also for wound healing¹³. From all these assessments, the medicinal benefits of *C. cainito* are supported.

Furthermore, the root of *C. cainito* has been utilized as alternative medicine in curing sterility, sexual asthenia, and

asthma; while seeds were predominantly used to alleviate hemorrhoid and intestinal worms. The bark has been wielded for treating cough, icterus, yellow fever while the fruits are used to treat dental decay and avitaminosis. Moreover, the entire plant can potentially treat ulcer and varicella¹⁴.

To further scientifically assess the medicinal importance of *C. cainito* Linn, a simple test on its cytotoxicity potential is highly desirable. Accordingly, cytotoxicity potential can be assessed by brine shrimp lethality bioassay (BSLT). The Brine shrimp lethality (BSLT) assay is an indispensable technique on the initial assessment of bioactive compounds present in plant extract¹⁵. This assay has been effective as a bioassay template in prescreening of active cytotoxic and antitumor agents¹⁶. Moreover, brine shrimp lethality assay provides a comprehensive analysis on the degree of cytotoxicity. It is a simple test with no aseptic techniques required. It can easily process a large quantity of organisms for statistical validation with relatively minimal quantity of sample¹⁵.

The test uses *Artemia salina* (Leach) reaction towards the extracts. The mortality is quantified and the lethality assay is computed. The assay is highly capable to detect wide spectrum of bioactivity in crude extracts. The method has been proven to be predictive of cytotoxicity and pesticide activity¹⁷. Previously, the brine shrimp lethality assay has been utilized in determining the cytotoxicity potential of various medicinal plants such as *Lantana camara*, *Chromolaena odorata*, and *Euphorbia hirta*¹⁸, *Kleinhovia hospita*¹⁹, *Acmella grandiflora*²⁰, *Ficus nota*²¹, *Phyllanthus niruri* and *Passiflora foetida*²².

In this study, the *C. cainito* Linn leaves were collected in Iligan City and their decoction, hydro-ethanolic, and ethanolic extracts were prepared and tested for their cytotoxicity potential against the brine shrimp and correlated with their known medicative properties of the plant. The data generated from the present work serve as baseline information in targeting specific bioactive compounds present in the *C. cainito* extracts.

Materials and methods

Botanical Source and Preparation of Extracts: *C. cainito* Linnaeus, particularly the leaves, was selected because of its known ethno-pharmacological uses based on a previous survey and interview with local folks and traditional practitioners in the communities. The fresh leaves of the plant were collected on February 2014 from a local source in Iligan City. The extracts from *C. cainito* leaves were prepared with various extraction solvent namely: ethanol, hydro-ethanol (50:50) and decoction. The decoction were prepared by cutting clean and fresh plant leaves into miniature sizes and heat to boil into distilled water in a 1:2 ratios for a minimum of 5 minutes. Plant samples were gradually chilled at 25°C before subjected to the process of separation then allow being freeze-dried succinctly to dissipate excess amount of moisture. In preparing the mixture of ethanolic extracts and hydro-ethanol, unwilt samples were

cleaned in clean water and then removed the dirt and other foreign matter by final rinsing using the sterile water. The cleaned samples were air-dried in a week or until the samples were already fragile upon picking. The dried samples were ground using a sterile electric blender. The powdered plant samples were weighed, and divided evenly into two parts and placed into a glass container; it was then filtered with sufficient quantity of absolute ethanol while the other samples was immersed in a 50:50 mixture of water-ethanol for three consecutive days (72h). The prepared solutions were purified using the whatman filter paper and decanted in storage glassware. An ample quantity of the purified ethanol solution was transferred in to rotary evaporation to prepare the ethanolic extract. The blended hydro-ethanol extract was accumulated in a *vacuo* and afterwards freeze-dried to formulate the hydroethanolic extract.

Brine Shrimp Lethality Test: The eggs of the Brine shrimp test organism eggs were acquired from MSU-IIT Chemistry Department. The sterile purified seawater was decanted in a devise compartment which simulates the light and dark areas. The eggs of the test organism were exposed into the dark portion of the compartment while the lamp above on the other boundary of the compartment captured the attention of the hatched shrimps. The larva of brine shrimps was employ for the cytotoxicity bioassay right after two days. From the three extracts (decoction, blended extract of hydro-ethanolic and ethanolic extract) with four concentrations of *C. cainito* were prepared: 10µg/mL, 100µg/mL, 500µg/mL and 1000µg/mL. By using the same amounts of the three extracts, the stock solutions were prepared. Then, 36.5mg, 25.18mg and 35.4mg were weighed and separately dissolved with plenty of solvent to prepare the 10,000-ppm stock solution. For the extraction of alcohol-based extracts, ethanol was used as extraction solvent and standby to be evaporated for two days. The same amounts of Dimethyl sulfoxide (DMSO) were added to the extracts except for the decoction. From the prepared stock solutions, 1000ppm, 500ppm, 100ppm and 10ppm concentrations from the stock solution were prepared by serial dilutions. For each extract, three replicates were prepared while the 5mL sterile filtered seawater served as the control set up. The Ten nauplii were added to every prepared extracts whereas another 10 nauplii were added to the control set up (sterile seawater). The test tubes were then observed and examined the number of dead (non-motile) nauplii in each test tube and was documented after 6 hours and 24 hours.

Data analysis: To analyze the result, the Reed-Muench statistical method was employed to assess the degree of toxicity of the *C. cainito* extracts to the test organisms. The response of *A. salina* was tested under various concentrations of the extract. The Lethality concentration (LC₅₀) constitutes the dose lethal to the 50 % of the population of the *A. salina*. To determine the lethality, the number of observed mortality (y-axis) was plotted versus log of concentration (x-axis). The concentration that inhibited 50% mortality constitutes the LC₅₀.

Results and discussion

The results showed that extraction with absolute ethanol and 50:50 ethanol-water successfully extracted the bioactive compounds in the *C. cainito* leaves. The effects of the various levels of concentrations of *C. cainito* Linnaeus extracts on the mortality of brine shrimp Nauplii were shown in Table-1. The brine shrimp mortality rates treated with the ethanolic and blended extract of hydro-ethanol was both 16.67% at 10 µg/ml and 100% at 1000 µg/ml. Meanwhile, the decoction extracts only brought about 16.67% and 96.67% mortality rates at 10 and 100 µg/ml, respectively. Consequently, the LC₅₀ range of the three extracts was 25.85 to 252 µg/ml. Based on the pattern of mortality rates, it can be inferred that the cytotoxic property of *C. cainito* is highly dose-dependent, as the concentration of the extract increased, the percentage of mortality rates and cytotoxicity of the extract towards the brine shrimp also increases. The cytotoxicity property of the extracts can be assessed based on the activity of the crude extract, it is highly toxic (active) if the value of LC₅₀ has less than 1000 ppm while non-toxic (inactive) if it is greater than 1000 ppm¹⁶. The results indicated that the three prepared extracts of *C. cainito* leaves have shown potential cytotoxic activity against brine shrimp.

It was previously reported that decocted leaves of *C. cainito* is used to treat various diseases of digestive systems such as

constipation, diarrhea, stomach ulcer, and rectal inflammation²³. The pharmacological properties of *C. cainito* may be rooted on its rich phytochemical contents such as flavonoids, anthraquinones, triterpenoids and notable elevated content of phenols^{24, 25}.

C. cainito has been reported to possess antioxidant properties²⁵ and wide range of antibacterial activities as it inhibits the growth of *Staphylococcus aureus*, *Micrococcus varians*, *Escherichia coli*, *Pseudomonas aeruginosa*, and *Proteus vulgaris*²⁶. Moreover, the ethanolic extract of *C. cainito* leaves was shown to have an antimicrobial activity against *S. aureus*, *E. coli*, *Salmonella typhimurium* and *Shigella spp.*²⁷.

It is highly plausible that the, the antimicrobial property of *C. Cainito* could be attributed to the ability to bind to the bacterial cell wall, thereby blocking its synthesis probably because of the saponins, flavonoids, tannin, steroid, and cardiac glycoside⁷. The other species of *Chrysophyllum* like the seed extracts of *C. albidum* possesses rich phytochemicals such as carbohydrates, cardiac glycosides, fatty acids, flavonoids saponins, quinones, and terpenoids while the aqueous extracts of the fruit of *C. albidum* possess antioxidant properties attributed to its high phenolic compounds²⁶.

Table-1: The effects of the various concentrations of *C. cainito* Linnaeus extracts on the mortality of brine shrimp Nauplii.

Extracts	Concentration, (µg/ml)	Mortality of brine Shrimp (%)		LC ₅₀ of Extract, (µg/ml)	
		After 6 H	After 24 H	Acute	Chronic
Ethanolic	1000	100	100	51.58	25.85
	500	86.67	100		
	100	76.67	93.33		
	10	16.67	16.67		
Hydrdo-ethanolic	1000	90	100	230.41	84.14
	500	100	100		
	100	16.67	53.33		
	10	0	16.67		
Decoction	1000	63.33	96.67	578.76	252.64
	500	53.33	73.33		
	100	6.67	23.33		
	10	10	16.67		

Moreover, the bark of other species of *Chrysophyllum*, the bark of *C. pruniforme* is also abundant with phytochemicals such as flavonoids, saponins, tannins, reducing sugars, polyphenols, and anthraquinones²⁸. So the cytotoxic activity of the ethanolic extracts can be attributed to its rich content of phytochemicals, antioxidants, and antimicrobial activities. The brine shrimp lethality assay proved to be a useful tool in the initial screening of potential bioactive compounds present in the plants. Moreover, to fully utilize the medicinal importance of *Chrysophyllum cainito*, further studies are highly desirable.

Conclusion

The result of the study demonstrated the toxicity of the ethanolic extract of *C. cainito*, which is very useful in the utilization of the species for further studies. The results indicated the presence of bioactive compounds which can be attributed to the plant's toxicological effects. Moreover, the results support the use of this plant species to be subjected for further pharmacological study to explore its specific medicinal properties. In addition, the present study supports the use of brine shrimp (*Artemia salina*) bioassay as a reliable, simple, and convenient method in the initial screening of bioactive compounds in medicinal plants.

References

1. Boerma, T. Mathers, C. AbouZahr, C. Chatterji, S. Daniel Hogan, D. and Gretchen Stevens (2020). Health in 2015 from Millenium Development goals (MDG'S) to Sustainable Developmentgoals (SDG'S). https://apps.who.int/iris/bitstream/handle/10665/200009/9789241565110_eng.pdf;jsessionid=9B66AE28DA01F191520B528067B5DC10?sequence=1, June 2020.
2. Debas H.T., Laxminarayan R. Straus SE. (2006). Complementary and alternative medicine. Chapter 69. In: Jamison DT, Breman JG, Measham AR, et al., editors. Disease control priorities in developing countries. 2nd ed. Washington DC: World Bank, 2006.
3. Ji, H.F. Li, X.J, Zhang H.Y. (2009). Natural products and drug discovery. EMBO Reports 10 (3), 194–200
4. Doan, H.V, and Le T.P. (2020). *Chrysophyllum cainito*: A Tropical Fruit with Multiple Health Benefits. Evidence-Based Complementary and Alternative Medicine 2020, 1-9. <https://doi.org/10.1155/2020/7259267>.
5. Rymbai, H., Dutta, S.K., Jha, A.K., Deshmukh, N.A., Verma, V.K., War, G.G., Paul, D., Patel, R.S., Chaudri, P., Mishra, L.K., Roy, A.R. and Roy, D. (2015). Chapter 32. Star Apple (*Chrysophyllum cainito*). https://www.researchgate.net/publication/335240511_STAR_APPLE_C_hrysophyllum_cainito/figures?lo=1&utm_source=google&utm_medium=organic, June 2020
6. Luo, X.D., Basile, M.J. and Kennelly, E.J. (2002). Polyphenolic antioxidants from the fruits of *Chrysophyllum cainito* (star apple). *J Agric Food Chem*, 50(6), 1379-1382.
7. Oranusi, S.U., Braide, W., Umeze, R.U. (2015). Antimicrobial activities and chemical compositions of *Chrysophyllum cainito* (star apple) fruit. *Microbiology Research International*, 3(3), 41-50.
8. Shailajan, S., and Gurjar, D. (2014). Pharmacognostic and Phytochemical Evaluation of *Chrysophyllum cainito* Linnaeus Leaves. *Int. J. Pharm. Sci. Rev. Res.*, 26(1), 106-111.
9. Hegde, K., Arathi, A., and Mathew, A. (2016). Evaluation of antidiabetic activity of hydro alcoholic extract of *Chrysophyllum cainito* fruits. *International Journal of Pharmaceutical Sciences and Research*, 7, 4422–4428.
10. Da Rosa, R.L., De Almeida, C.L., Somensi, L.B., Boeing, T., Mariano, L.N.B., Krueger, C.M.A., De Souza, P., Filho, V.C., Da Silva, L.M., and De Andrade, S.F. (2019). *Chrysophyllum cainito* (apple-star): A fruit with gastroprotective activity in experimental ulcer models. *Inflammopharmacology* 27(5), 985-996. <http://doi:10.1007/s10787-017-0427-z>
11. Ernest, A. K., Marie-Solange, T., Beugre, K., & Noeuml, Z. G. (2009). Effect of aqueous extract of *Chrysophyllum cainito* leaves on the glycaemia of diabetic rabbits. *African Journal of Pharmacy and Pharmacology*, 3(10), 501-506.
12. Meira N.A., Klein, L.C., Rocha, L.W., Quintal, Z.M., Monache, F.D., Filho, V.C. and Quintão, N.L.M. (2014). Anti-inflammatory and anti-hypersensitive effects of the crude extract, fractions and triterpenes obtained from *Chrysophyllum cainito* leaves in mice. *Journal of Ethnopharmacology*, 151(2), 975-983. <http://doi:10.1016/j.jep.2013.12.014>
13. Shailajan, S., and Gurjar, D. (2016). Wound healing activity of *Chrysophyllum cainito* L. leaves: evaluation in rats using excision wound model. *Journal of Young Pharmacists*, 8(2), 96–103.
14. Houessou, L.G., Loughbegnon, T.O., Gbesso, F.G.H., Anagonou, L.E.S., Sinsin, B. (2012). Ethno-botanical study of the African star apple (*Chrysophyllum albidum* G. Don) in the Southern Benin (West Africa). *Journal of Ethnobiology and Ethnomedicine*, 8, 1–10
15. Sarah, Q.S., Anny, F.C., and Misbahuddin, M. (2017). Brine shrimp lethality assay. *Bangladesh Journal Pharmacology*, 12, 186-189.
16. Meyer, N., Ferrigni, R., Putnam, E., Jacobsen, B., Nichols, E., and McLaughlin, L. (1982). Brine Shrimp: A convenient General Bioassay for Active Plant Constituents. *Hippocrates Verlag GmbH.*, 45, 31-34.
17. Mentor, R., Jovanova, B. and Panovska, T.K. (2014). Toxicological evaluation of the plant products using Brine Shrimp (*Artemia salina* L.) model. *Macedonian Pharmaceutical Bulletin*, 60(1), 9-18.

18. Olowa, L. and Nuñez, O. (2013). Brine Shrimp Lethality Assay of the Ethanolic Extracts of Three Selected Species of Medicinal Plants from Iligan City, Philippines. *International Research Journal of Biological Sciences*, 2(11), 74-77.
19. Morilla, L.J., Nuñez, O. and Uy, M. (2015). Brine Shrimp Lethality Test of *Kleinhovia hospita* stem and bark from Agusan del Sur, Philippines. *ELBA Bioflux*, 7(1).
20. Elias, N., Nuñez, O. and Uy, M. (2014). Evaluating the Potential Cytotoxic Activity of *Acmella grandiflora* Flower and Whole Plant using Brine Shrimp Lethality Test. *International Research Journal of Biological Sciences*, 3(10), 90-92.
21. Arquion, C., Nuñez, O. and Uy, M. (2015). Evaluating the Potential Cytotoxic Activity of *Ficus nota* Using Brine Shrimp Lethality Test. *Bulletin of Environment, Pharmacology and Life Sciences*, 4(12).
22. Juario, J., Mondejar, E., Nuñez, O. and Uy, M. (2015). Effect of *Phyllanthus niruri* and *Passiflora foetida* Extracts on the Mortality and Survival rate of the Brine Shrimp *Artemia salina*. *Research Journal of Recent Sciences*, 4(2), 61-67.
23. Balinado, L. and Chan, M. (2017). An Ethnomedicinal Study of Plants and Traditional Health Care Practices in District 7, Cavite, Philippines. *International Conference on Chemical, Agricultural, Biological and Medical Sciences (CABMS-17)*.
24. Chel-Guerrero, L., Sauri-Duch, E., Fragoso-Serrano, M.C., Pe'rez-Flores, L.J., Go'mez-Olivares, J.L., Salinas-Arreortua, N., Sierra-Palacios, E., and Mendoza-Espinoza, J.A. (2017). Phytochemical Profile, Toxicity, and Pharmacological Potential of Peels from four Species of Tropical Fruits. *Journal of Medicinal Food*, 00(0), 1-10.
25. Li, L.B., Lin, S., Yan, J., Wang, Q.L., Fan, Z.Y., Dong, Q.R., Qin, J.Z., and Xie, Z.G. (2015). Poly-phenolic fraction of *Chrysophyllum cainito* extract induces cell death in osteosarcoma cells. *Bangladesh Journal of Pharmacology*, 10, 972-979.
26. Oputah, S., Mordi, R., Ajanaku, K., Olugbuyiro, J., Shade, J., Olorunshola, O., and Azuh, D. (2016). Phytochemical and Antibacterial Properties of Ethanolic Seed Extracts of *Chrysophyllum albidum* (African Star Apple). *Oriental Journal of Physical Sciences*, 1(1 & 2), 05-09.
27. Duyilemi, O.P., and Lawal, I.O. (2009). Antibacterial activity and phytochemical screening of *Chrysophyllum albidum* leaves. *Asian Journal of Food and Agro-Industry*, S75-S79.
28. Angone, S., Mewono, L., Boukandou, M.M., Medzegue, M.S., Ndong, J., Djoba Siawaya, J.F. and Souza, A. (2013). Phytochemical screening and cytotoxicity studies of *Chrysophyllum pruniforme* Pierre ex Engl. Barks. *Pharmacognosy Research*, 5(3).